

Graph Covers


Aleksander Malnič
University of Ljubljana

Niš, Srbija

November 10, 2009


Covering projections of graphs

Covering projections of graphs



A surjective mapping $p: \tilde{X} \rightarrow X$ such that $D_{\tilde{u}} \rightarrow D_u$ is a bijection

Covering projections of graphs



A surjective mapping $p: \tilde{X} \rightarrow X$ such that $D_{\tilde{u}} \rightarrow D_u$ is a bijection

\tilde{X} : covering graph, X : base graph

$p^{-1}(u), p^{-1}(x)$: fibres

Regular cover : fibres are orbits of a semiregular $\text{CT}(p) \leq \text{Aut}\tilde{X}$

1. Complete solution of the long-standing Heawood's Map Colour Problem by Ringel and Youngs 74'. Use of covers first recognized by Alpert and Gross 74'.

Motivation I

1. Complete solution of the long-standing Heawood's Map Colour Problem by Ringel and Youngs 74'. Use of covers first recognized by Alpert and Gross 74'.
2. Combinatorial treatment of graph covers by Gross 74'. Same idea in Biggs' monograph Algebraic Graph Theory 74' (in connection with symmetries of graphs).

1. Complete solution of the long-standing Heawood's Map Colour Problem by Ringel and Youngs 74'. Use of covers first recognized by Alpert and Gross 74'.
2. Combinatorial treatment of graph covers by Gross 74'. Same idea in Biggs' monograph Algebraic Graph Theory 74' (in connection with symmetries of graphs).
3. Systematic combinatorial theory of graph covers by Gross and Tucker 74'-77', Topological graph Theory 87'. Extended to graph bundles by Pisanski and Vrabec 82'.

1. Complete solution of the long-standing Heawood's Map Colour Problem by Ringel and Youngs 74'. Use of covers first recognized by Alpert and Gross 74'.
2. Combinatorial treatment of graph covers by Gross 74'. Same idea in Biggs' monograph Algebraic Graph Theory 74' (in connection with symmetries of graphs).
3. Systematic combinatorial theory of graph covers by Gross and Tucker 74'-77', Topological graph Theory 87'. Extended to graph bundles by Pisanski and Vrabec 82'.
4. Systematic combinatorial theory of branched coverings of surfaces by Gross and Tucker 74'-77', Topological graph Theory 87'. Also by Jones and Singerman 78'.

Motivation II - studying symmetries of graphs

Motivation II - studying symmetries of graphs

5. Vertex, edge, arc transitive, of small valencies

Conjecture of Marušič: every vertex transitive graph is a regular \mathbb{Z}_p -cover of some smaller graph. Confirmed for valencies 3 and 4.

Motivation II - studying symmetries of graphs

5. Vertex, edge, arc transitive, of small valencies

Conjecture of Marušič: every vertex transitive graph is a regular \mathbb{Z}_p -cover of some smaller graph. Confirmed for valencies 3 and 4.

5.1 Compiling lists of highly symmetric graphs

Foster census (Bouwer '88): arc-trans. cubic graphs on ≤ 512 vertices (lattice of regular covers). Extended to 768 vertices by Conder and Dobcsányi 02'. List of semisymmetric cubic graph ≤ 768 vertices by Conder, M, Marušič, Potočnik 03'.

Motivation II - studying symmetries of graphs

5. Vertex, edge, arc transitive, of small valencies

Conjecture of Marušič: every vertex transitive graph is a regular \mathbb{Z}_p -cover of some smaller graph. Confirmed for valencies 3 and 4.

5.1 Compiling lists of highly symmetric graphs

Foster census (Bouwer '88): arc-trans. cubic graphs on ≤ 512 vertices (lattice of regular covers). Extended to 768 vertices by Conder and Dobcsányi 02'. List of semisymmetric cubic graph ≤ 768 vertices by Conder, M, Marušič, Potočnik 03'.

5.2 Construction of infinite families

Tutte 46': cubic graphs ≤ 5 -arc trans. Few 5-arc trans. examples known. First infinite family by Djoković 74' (reg. covers of Tutte's 8-cage).

Motivation II - studying symmetries of graphs

5. Vertex, edge, arc transitive, of small valencies

Conjecture of Marušič: every vertex transitive graph is a regular \mathbb{Z}_p -cover of some smaller graph. Confirmed for valencies 3 and 4.

5.1 Compiling lists of highly symmetric graphs

Foster census (Bouwer '88): arc-trans. cubic graphs on ≤ 512 vertices (lattice of regular covers). Extended to 768 vertices by Conder and Dobcsányi 02'. List of semisymmetric cubic graph ≤ 768 vertices by Conder, M, Marušič, Potočnik 03'.

5.2 Construction of infinite families

Tutte 46': cubic graphs ≤ 5 -arc trans. Few 5-arc trans. examples known. First infinite family by Djoković 74' (reg. covers of Tutte's 8-cage).

5.3 Classifying graphs

Trans. graphs via orbital graphs (essentially: a trans. perm. group and its stabilizer). Djoković and Miller 80' classified stabilizers of arc trans. cubic graphs. Extended to edge-trans. cubic graphs by Goldschmidt 80'.

Praeger's normal reduction

Praeger's normal reduction

THM, Praeger 88'

$\text{Aut} X$ 2-arc trans, $N \triangleleft \text{Aut} X$ with ≥ 3 orbits. Then $X \rightarrow X/N$ is a regular cover, with $\text{Aut} X/N$ acting 2-arc trans. on X/N .

Praeger's normal reduction

THM, Praeger 88'

$\text{Aut}X$ 2-arc trans, $N \triangleleft \text{Aut}X$ with ≥ 3 orbits. Then $X \rightarrow X/N$ is a regular cover, with $\text{Aut}X/N$ acting 2-arc trans. on X/N .

Base graphs: every normal subgroup has ≤ 2 orbits.

Praeger's normal reduction

THM, Praeger 88'

$\text{Aut}X$ 2-arc trans, $N \triangleleft \text{Aut}X$ with ≥ 3 orbits. Then $X \rightarrow X/N$ is a regular cover, with $\text{Aut}X/N$ acting 2-arc trans. on X/N .

Base graphs: every normal subgroup has ≤ 2 orbits.

Classify base graphs (usually using Classification of finite simple groups). The rest are normal covers. Can we give a neat description?

Key idea: Lifting automorphisms

Key idea: Lifting automorphisms

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\sigma^z} & \tilde{X} \\ p \downarrow & & \downarrow p \\ X & \xrightarrow{\sigma^q} & X. \end{array}$$

Key idea: Lifting automorphisms

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\sigma^{\tilde{z}}} & \tilde{X} \\ p \downarrow & & \downarrow p \\ X & \xrightarrow{\sigma^g} & X. \end{array}$$

All lifts of $G \leq \text{Aut} X$ constitute a group $\tilde{G} \leq \text{Aut} \tilde{X}$. There is a short exact sequence $1 \rightarrow \text{CT}(p) \rightarrow \tilde{G} \rightarrow G \rightarrow 1$, and any abstract group extension can be studied as a lifting problem.

Key idea: Lifting automorphisms

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\sigma\tilde{z}} & \tilde{X} \\ p \downarrow & & \downarrow p \\ X & \xrightarrow{\sigma} & X. \end{array}$$

All lifts of $G \leq \text{Aut}X$ constitute a group $\tilde{G} \leq \text{Aut}\tilde{X}$. There is a short exact sequence $1 \rightarrow \text{CT}(p) \rightarrow \tilde{G} \rightarrow G \rightarrow 1$, and any abstract group extension can be studied as a lifting problem.

THM, Djoković 74'

Let $p: \tilde{X} \rightarrow X$ regular cover. If $G \leq \text{Aut}X$, s-trans., lifts, then \tilde{G} is s-trans.

Key idea: Lifting automorphisms

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\sigma\tilde{z}} & \tilde{X} \\ p \downarrow & & \downarrow p \\ X & \xrightarrow{\sigma} & X. \end{array}$$

All lifts of $G \leq \text{Aut}X$ constitute a group $\tilde{G} \leq \text{Aut}\tilde{X}$. There is a short exact sequence $1 \rightarrow \text{CT}(p) \rightarrow \tilde{G} \rightarrow G \rightarrow 1$, and any abstract group extension can be studied as a lifting problem.

THM, Djoković 74'


Let $p: \tilde{X} \rightarrow X$ regular cover. If $G \leq \text{Aut}X$, s-trans., lifts, then \tilde{G} is s-trans.

Djoković used topological approach via fundamental groups. Not appropriate to cope with specific problems arising with graphs.


Covers and Lifting automorphisms combinatorially

Covers and Lifting automorphisms combinatorially



Regular covers by regular voltages in $\Gamma \cong \text{CT}(p)$



Z_2^3




Z_2




Covers and Lifting automorphisms combinatorially



Regular covers by regular voltages in $\Gamma \cong \text{CT}(p)$



\mathbb{Z}_2^3



\mathbb{Z}_2



Basic lifting lemma (for regular covers)

$g \in \text{Aut} X$ lifts \Leftrightarrow the mapping $\text{vol}(W) \rightarrow \text{vol}(W^g)$ of voltages of the fundamental closed walks at some vertex extends to an automor. of Γ

Abelian regular covers

Abelian regular covers

G has a matrix representation G_{\sharp} on $H_1(X, \mathbb{Z})$. Consider the transposed matrix group G_{\sharp}^t . Then

Abelian regular covers

G has a matrix representation G_{\sharp} on $H_1(X, \mathbb{Z})$. Consider the transposed matrix group G_{\sharp}^t . Then

THM., M., Marušič, Potočnik, 03'

$G \in \text{Aut} X$ lifts \Leftrightarrow the base homology cycles of X are given voltages which represent a basis of an G_{\sharp}^t -invariant submodule of $H_1(X, \mathbb{Z})$.

Abelian regular covers

G has a matrix representation G_{\sharp} on $H_1(X, \mathbb{Z})$. Consider the transposed matrix group G_{\sharp}^t . Then

THM., M., Marušič, Potočnik, 03'

$G \in \text{Aut} X$ lifts \Leftrightarrow the base homology cycles of X are given voltages which represent a basis of an G_{\sharp}^t -invariant submodule of $H_1(X, \mathbb{Z})$.

Invariant submodules determine the covers up to equivalence. There is a way to reduce the obtained covers up to isomorphism.

Abelian regular covers

G has a matrix representation G_{\sharp} on $H_1(X, \mathbb{Z})$. Consider the transposed matrix group G_{\sharp}^t . Then

THM., M., Marušič, Potočnik, 03'

$G \in \text{Aut} X$ lifts \Leftrightarrow the base homology cycles of X are given voltages which represent a basis of an G_{\sharp}^t -invariant submodule of $H_1(X, \mathbb{Z})$.

Invariant submodules determine the covers up to equivalence. There is a way to reduce the obtained covers up to isomorphism.

Particularly useful when dealing with elementary abelian \mathbb{Z}_p^k -covers. We need to find invariant subspaces of matrix groups acting on \mathbb{Z}_p^r , where $r = \text{betti}(X)$.

Abelian regular covers

G has a matrix representation G_{\sharp} on $H_1(X, \mathbb{Z})$. Consider the transposed matrix group G_{\sharp}^t . Then

THM., M., Marušič, Potočnik, 03'

$G \in \text{Aut} X$ lifts \Leftrightarrow the base homology cycles of X are given voltages which represent a basis of an G_{\sharp}^t -invariant submodule of $H_1(X, \mathbb{Z})$.

Invariant submodules determine the covers up to equivalence. There is a way to reduce the obtained covers up to isomorphism.

Particularly useful when dealing with elementary abelian \mathbb{Z}_p^k -covers. We need to find invariant subspaces of matrix groups acting on \mathbb{Z}_p^r , where $r = \text{betti}(X)$.

For concrete covers one can use computer base packages like Magma for fast computation of invariant subspaces (MeatAxe algorithm).

Example

Example

Find all regular elementary abelian covers of the 3-dipole such that some edge transitive subgroup lifts.

Example

Find all regular elementary abelian covers of the 3-dipole such that some edge transitive subgroup lifts.

Minimal edge transitive subgroup cyclically permutes the edges, fixing the two vertices. Induced action on base cycles is given by

$$R = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix} \quad R^t = \begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}$$

The full space gives rise to the homological cover. Proper nontrivial invariant subspaces are 1-dim. $\Delta = \lambda^2 + \lambda + 1$, and the eigenvalues depend on the congruence type of $p \pmod 3$.

$p = 3$, then $\lambda = 1$, and $\underline{v} = [1, 2]^t$

$p \not\equiv -1 \pmod 3$, then Δ irreducible

$p \equiv 1 \pmod 3$, then $\lambda = -\xi, -\xi^2$, and $\underline{v} = [1, -\xi^2]^t, [1, -\xi]^t$

The eigenvectors induce voltage assignments on arcs, and hence determine the derived graphs. The obtained covers are pairwise nonequivalent, but the last two are isomorphic.

Classifying transitive covers of intransitive graphs

Classifying transitive covers of intransitive graphs

We consider regular abelian covers, and modified version of Praeger's normal reduction. Take quotients $X \rightarrow X/N$ by normal subgroups arising from linear representation on eigenspaces of X .

Classifying transitive covers of intransitive graphs

We consider regular abelian covers, and modified version of Praeger's normal reduction. Take quotients $X \rightarrow X/N$ by normal subgroups arising from linear representation on eigenspaces of X .

These subgroups can be computed in terms of eigenvalues and eigenvectors of the base graph, using complex irreducible characters of abelian voltage groups.

Thank you!